

— 2018 —
WATER YEAR

ENVISIONING
BINATIONAL
COLLABORATION

a living river

CHARTING SANTA CRUZ RIVER CONDITIONS—NOGALES WASH TO AMADO
SUPPLEMENTARY REPORT FOR 2008 TO 2018 WATER YEARS



SONORAN
INSTITUTE

SANTA CRUZ RIVER A LIVING ECOSYSTEM


The Santa Cruz River in Santa Cruz County flows year round from Rio Rico to Amado and supports a culturally and ecologically diverse region. River flows are sustained by effluent discharges from the Nogales International Wastewater Treatment Plant (NIWTP) which treats wastewater from Nogales, Arizona and Nogales, Sonora (collectively *Ambos Nogales*). Keeping this stretch of the river flowing is critical to the viability of this ecologically rich and culturally significant area. With Mexico legally controlling over 80% of the water released into the river, any effort to permanently dedicate water to the river will require a binational solution that benefits both countries. In the last decade, two important changes in wastewater management have occurred:

Upgraded Treatment Plant: In 2009 the NIWTP completed significant upgrades to the treatment process resulting in reduced levels of nitrogen in the released effluent.

Wastewater Diversions: In 2013 Nogales, Sonora completed construction of the Los Alisos treatment plant to divert and treat some of the wastewater that would otherwise be sent to NIWTP. Effluent generated at Los Alisos is released into a different river and flows south into Sonora.

Rivers are dynamic with conditions influenced by many factors. How do we know if management actions and other factors are influencing the conditions of this living ecosystem? The *Living River* reports were developed to annually track indicators of river


conditions and determine what is changing. This assessment summarizes data from the 2008–2018 water years (each water year spans from October 1 to September 30). This year we include conditions in Nogales Wash, an important corridor linking Ambos Nogales to the NIWTP and to the river. The pages following this executive summary provide more detail on the water context and data for 12 indicators of river health.



CHANGES IN AQUATIC AND RIPARIAN CONDITIONS

- **Improved water quality with decreased nutrient pollution:** Nitrogen and phosphorus are essential nutrients for plants and animals, but too much can degrade conditions for aquatic life. Total phosphorus and ammonia, a form of nitrogen which can be toxic to fish, significantly decreased in the Santa Cruz after the treatment plant upgrades were complete.
- **Availability of dissolved oxygen increases:** Dissolved oxygen, essential for aquatic life, increased in the Santa Cruz after the treatment plant upgrades were complete.
- **Pollutant metals no longer threaten aquatic wildlife:** Cadmium, a metal that is lethal to wildlife at low concentrations, has not been detected since 2011.

- **More diverse aquatic life:** Improvements in water quality has allowed aquatic life in the river to rebound. Four species of fish, including the endangered Gila topminnow, are now found. Diversity of aquatic invertebrates has increased and more pollution-sensitive species are present.
- **Reduced flow extent:** The length of the flowing river has decreased, likely the result of a combination of factors including increased water infiltration with reduced nutrient pollution, scouring floods, and water diversions. Initial diversions to Los Alisos appear to be minor and reductions in effluent released into the river is primarily due to a decrease in wastewater from Nogales, Arizona.
- **Stressed cottonwoods at the far end of the river:** Branch die-back and death of some trees indicate that cottonwoods in the Amado Reach are suffering. Reduced flow extent is likely the primary cause of stress as flow has become more irregular through this reach.



OTHER OBSERVATIONS

- **Fecal contamination continues to be a risk during the rainy seasons:** *E. coli* levels continue to exceed the swimming (full body contact) standard. Exceedances of *E. coli* primarily occur during rainy seasons suggesting multiple sources of fecal contamination exist throughout

the watershed that likely come from humans, livestock, and wildlife.






- **Groundwater levels most stable in the Tubac Reach:** Both the maximum depth to groundwater and the annual January to June decline in groundwater levels are sufficiently shallow and stable to provide optimal conditions for cottonwoods in the Tubac Reach.

ASSESSING CONDITIONS

This Living River report evaluates conditions of the Santa Cruz River from Nogales Wash to Amado using indicators (see table below) organized into four categories that represent a breadth of biological, chemical, and physical properties of the river.

The indicators relate to conditions in the river channel and in the riparian areas, the areas next to and affected by the river.

The purpose of the *Living River* series is to monitor and report on river conditions in Nogales Wash and at various

CATEGORY		PURPOSE	INDICATORS
Flow Extent		Water flowing in and out of the system determines available aquatic habitat.	<ul style="list-style-type: none"> • Miles of flow in June • Number of dry days or visits
Water Quality		Specific chemical conditions are necessary to sustain the river's animal and plant communities.	<ul style="list-style-type: none"> • Dissolved oxygen • Total phosphorus • Ammonia • <i>E. coli</i> • Metals
Aquatic Wildlife		Wildlife in the river integrate and reflect conditions of many factors of the surrounding environment.	<ul style="list-style-type: none"> • Fish • Aquatic invertebrates
Groundwater		Mature riparian trees require certain surface and groundwater conditions to survive.	<ul style="list-style-type: none"> • Maximum depth • January to June decline
Riparian vegetation*		Plant communities reflect changes in water quantity and quality.	<ul style="list-style-type: none"> • Forest and woodland cover

intervals downstream of the effluent discharge point. As effluent flows downstream, it impacts and is impacted by the natural conditions of soils, vegetation, and the surrounding ecosystem. For the purposes of this study, the 20-mile stretch of Santa Cruz River is divided into three sections, or reaches: Rio Rico, Tubac, and Amado. These reaches differ in geology and hydrology, with portions of the reaches alternating between gaining and losing stream conditions. There is a close connection between the surface and groundwater that can influence the aquatic and riparian

ecosystems in these reaches. Most of the data for Nogales Wash come from a single survey location near Ruby Road, though occasionally data also come from a site closer to the international border.

Data are collected and summarized by water year (October 1–September 30) and compared to the baseline conditions observed in the 2008 water year. This supplemental report shares data from all the water years to enable an easy viewing of trends and long-term patterns.

*Riparian vegetation only monitored from 2008–2010


Streamflow and Rainfall

The amount of water flowing in the river provides important context when examining the results of indicators that track the conditions along the Santa Cruz from Nogales Wash to Amado. This water comes from two natural sources: precipitation and groundwater. A third source of water, effluent, is water that has been used by people, treated in a wastewater facility, and returned to a river. The Nogales International Wastewater Treatment Plant continuously releases effluent into Santa Cruz, just downstream of where the Nogales Wash flows into the river. Wastewater coming from Ambos Nogales flows in a pipeline, the International Outfall Interceptor (IOI), that is buried beneath Nogales Wash. Nogales Wash is also the principal waterway transporting stormwater runoff from Ambos Nogales to the river. Large, uncontrolled flows of stormwater, together with accidental sewage discharges from the IOI, have raised flood safety and water quality concerns from communities on both sides of the border.

Effluent accounts for the majority of streamflow in the Santa Cruz River from Rio Rico to Amado. However, streamflow also includes stormwater from the watershed. The Santa Cruz River watershed includes all of the land whose stormwater flows toward the river. We can get a sense of stormwater contribution to the Santa Cruz by tracking total rainfall and streamflow upstream of the treatment plant in Nogales Wash and near the international border. Seasonal floods (which can be measured by looking at peak flows) are important for scouring the riverbed, recharging aquifers, dispersing seeds, inducing seed germination, and clearing natural debris.

2008–2018 Summary

The total volume of effluent released annually into the river has varied, but has decreased by an average of 2,000 acre-feet (over 650 million gallons) since 2013. This represents about a 12% decrease compared to 2008. Although diversions of wastewater to the new Los Alisos Treatment



Streamflow and Rainfall cont.


plant in Nogales, Sonora began in 2013, this decrease in effluent volume primarily results from reduced volume of wastewater coming from Nogales, Arizona. Over time, there could be further reductions of effluent with increased diversions of wastewater to the Los Alisos treatment plant in Nogales, Sonora.

Streamflow is measured daily in Tubac, near the international border, and since 2011, along Nogales Wash. Overall, total volume of streamflow at the Tubac gage has decreased since 2008. The 2017 water year is the most recent peak in total volume of water passing this gage, largely because of high flows during the summer monsoon.

Total rainfall has hovered around the recent average at Nogales Airport. Sixteen miles to the northwest, rainfall near Tumacácori National Historical Park has been lower than the historical average in this area.

Peak streamflow has varied with location. Flooding conditions have generally been higher upstream of the treatment plant than in Tubac with the largest flood recorded at the Nogales gage near the international border. Flows in Tubac have been more variable, with the highest peak flow occurring in 2014. Nogales Wash has seen an increase in peak flow volume that has been fairly stable since 2014.

With overall effluent volume remaining fairly constant, the decrease in total volume of stream flow could be due to several factors including changing weather patterns, floods, variability in groundwater levels, and reduced nutrient pollution. Cleaner water promotes infiltration of water through the riverbed by breaking down old clogging layers, preventing new layer from forming, and ultimately recharging the aquifer.






FLOW EXTENT: Miles of Flow

Measuring flow extent, or the distance the river is flowing, is a quick visual technique to track changes in the amount of water in the river and provides a rough estimate of the quantity of aquatic habitat available. For example, high flow extent may indicate availability of habitat for aquatic life. Low flow extent may indicate reduced water inputs, which could decrease aquatic habitat. Alternatively, low flow extent could indicate greater recharge of water into local aquifers.

Measuring miles of flow prior to the monsoon season determines the minimum extent of flow during the driest time of year. This is typically measured in mid-June. Friends of the Santa Cruz River track river conditions monthly at three sites along the river and streamflow is monitored continuously by a USGS stream gage in Tubac. While the exact location of the "end of flow" is unknown, we can track approximate location by knowing which monitoring sites are dry at the time of survey in June. Flow extent is not measured in Nogales Wash.

2008–2018 Summary

Miles of flow in June has decreased since the 2008 baseline. Prior to 2010, the river flowed 17 miles and into the Amado reach regularly in June. Since the volume of effluent released into the river did not significantly decrease after the 2009 upgrade, increased recharge from improved water quality is likely a major factor for the decreased flow extent in 2010 and 2011. A further reduction in miles of flow was observed in 2013 and 2014. This coincides with the timing of diversions of wastewater to a new wastewater facility in Sonora, Mexico which began in 2013. However, the total volume of effluent released into the river has decreased only by an average of 2,000 acre-feet per year, with this decrease largely attributed to decreased wastewater coming from Nogales, Arizona. Since 2014, flow extent in June has increased and stabilized.





FLOW EXTENT: Number of Dry Days or Visits

In addition to measuring minimum flow extent in the driest part of the year, it is helpful to understand the variation throughout the year. Although we can't know the exact length of the river at all times, quantifying the amount of time a specific location along the river is dry gives a measure of change in flow extent across the year.

Friends of the Santa Cruz River track river conditions monthly at three sites along the river and one site on Nogales Wash. Streamflow is monitored continuously by a stream gage in Tubac managed by the U.S. Geological Survey. Measuring the number of monthly visits or days that are dry at a given location provides an estimate of changes in amount of water in the river.

2008–2018 Summary


Dry days or visits have become more common at the far downstream end of the reach, indicating that flow extent has

decreased. Prior to 2010, the river flowed every day at the Tubac stream gage with no zero flow days recorded since July of 2003. Since 2010, the number of dry days has increased, peaking at nearly 120 in the 2014 water year.

A similar pattern of increased dryness is seen at the start of the Amado reach. Though measured monthly rather than daily, the number of dry visits has increased. Only one dry visit was observed between 2005 and 2009. In contrast, dry visits peaked during the 2013 and 2014 water years with 9 of 12 monthly visits dry. Since 2014, dry days and visits have decreased and stabilized.

Nogales Wash has had stable flows. Only three of the monthly monitoring visits at Nogales Wash found no flow. These were during the June visits of 2011, 2013, and 2014.

Direction of Flow >>>>



Note: Data from the Rio Rico reach are not displayed. No dry visits observed during the 2008–2018 water years.



WATER QUALITY: Dissolved Oxygen


Fish and other aquatic animals need oxygen to survive. Rivers absorb oxygen from the atmosphere and aquatic plants produce oxygen. External causes of variability in dissolved oxygen levels include nutrient levels, groundwater discharge, shading, water temperature, and time of day.

The Arizona Department of Environmental Quality sets the minimum wildlife standard for dissolved oxygen in effluent-dependent streams at 1 milligram per liter (mg/L) measured between sunset and three hours after sunrise, and 3 mg/L at other times. All data were measured within three hours after sunrise, thus measures of dissolved oxygen greater than 1 mg/L meet the standard.

2008–2018 Summary

Levels of dissolved oxygen have met the standard in all three reaches of the Santa Cruz. Dissolved oxygen increased after upgrades to the treatment plant were completed in 2009. Although there have been more days with no flow in the Amado reach in recent years, when water is present, levels of dissolved oxygen have been high.

In Nogales Wash, the standard for dissolved oxygen is less stringent because this is a warm water system that isn't dependent on effluent releases for flow. Levels of dissolved oxygen have been stable and 92 of 134 samples have met the standard, or they attain the standard 69% of the time. Treatment plant upgrades did not impact conditions in Nogales Wash.





WATER QUALITY: Total Phosphorus


Phosphorus is an essential nutrient for plant and animal life. However, too much phosphorus can reduce the quality of aquatic habitat. Elevated levels of phosphorus can accelerate plant growth. As plants decay in the river, dissolved oxygen decreases and aquatic animals can suffocate.

The median amount of total phosphorus monitored in the Rio Rico reach from 1992-1999 was approximately 5 mg/L. Because Arizona Department of Environmental Quality does not have a standard for total phosphorus in this stretch of the Santa Cruz River, this assessment uses the 1992-1999 median of 5 mg/L as the historical standard. Similarly, there is no standard for phosphorus in Nogales Wash. This assessment uses the 1996-1999 median of <1 mg/L as a historical standard.

2008-2018 Summary

Phosphorus levels declined over the years in the Rio Rico and Tubac reaches, most notably after the upgrades to the treatment plant were complete in 2009. Data are limited for the Amado reach but measures of total phosphorus are below the historical standard.

Nogales Wash has limited data and treatment plant upgrades did not impact conditions. All but five samples met the historical standard. The only sample since 2013 was taken in 2017 during a sewage breach when untreated wastewater was flowing out of the International Outfall Interceptor and into Nogales Wash.





WATER QUALITY: Ammonia


Nitrogen is an essential nutrient for plant and animal life, but too much can lead to nutrient pollution. Ammonia (NH_3) is one form of nitrogen that can be toxic to fish. Even at low concentrations, ammonia can reduce hatching success, among other impacts. The toxicity of ammonia varies with several factors such as pH, temperature, and dissolved oxygen. The Arizona Department of Environmental Quality's wildlife standard for ammonia varies by pH and temperature. As pH and temperature increase, the toxicity of ammonia increases, thus the acceptable level of ammonia decreases with high pH and temperature.

High nutrient levels can also increase the number of microorganisms that break down and use these nutrients. Living in the spaces between the sand and gravel of the streambed, these organisms can become so numerous that they create a clogging layer and prevent water from moving through the streambed to recharge the local aquifer.

2008–2018 Summary

Ammonia decreased dramatically and was often not detected in the Rio Rico and Tubac reaches after the treatment plant upgrade was complete in 2009. This improved water quality led to beneficial conditions for fish and other aquatic wildlife. Although data are limited for the Amado reach during this period, between 2000 and 2007 measures of ammonia averaged 10 mg/L and followed a similar pattern to the Rio Rico and Tubac reaches. The decreased ammonia likely reduced the clogging effect in the river bed and was a factor in the reduced flow extent observed since 2009.

Nogales Wash was not impacted by the upgrades to the treatment plant. Ammonia levels here were low, averaging 1.4 mg/L from 2008–2013. Measures of ammonia from 2017 were taken during a sewage breach when untreated wastewater was flowing out of the International Outfall Interceptor and into Nogales Wash.



Note - Ammonia standards vary with temperature and pH and can't be graphed as a single line.



WATER QUALITY: *E. coli*


Escherichia coli (*E. coli*) is one of the common species of bacteria living in the lower intestines of mammals, and its presence in water is an indication of fecal contamination. The discovery of *E. coli* indicates the potential presence of other pathogenic microorganisms such as bacteria and viruses that might be a health risk to people swimming or wading in the river.

The Arizona Department of Environmental Quality standard for a single sample maximum for full body contact (swimming) is 235 colony-forming units per 100 milliliters of water (CFU/100mL). For partial body contact (wading), the single sample maximum is 576 CFU/100mL. The results are compared to the stricter standard, thus samples with greater than 235 CFU/100mL do not meet the standard.

2008–2018 Summary

E. coli levels varied seasonally and were similar in all three reaches of the river and in Nogales Wash. The Nogales International Wastewater Treatment Plant has always removed bacteria prior to release of effluent in the Santa Cruz, thus levels of *E. coli* in the effluent did not change after the upgrades were complete. Levels of *E. coli* exceeded the standard most often during the rainy seasons. High levels during the rainy season suggest that rain is washing fecal contamination into the river from multiple sources within the watershed. Preliminary research identifies humans, livestock, and wildlife among the sources of contamination.

Several of the exceedances in Nogales Wash from 2017 were from samples taken during a sewage breach when untreated wastewater was flowing out of the International Outfall Interceptor and into Nogales Wash.





WATER QUALITY: Metals

Metals in high concentrations endanger wildlife in aquatic ecosystems by lowering reproductive success, interfering with growth and development, and, in extreme cases, causing death. Most metals build up in aquatic food chains and may pose long-term threats to all organisms in the aquatic environment. The Santa Cruz River is exposed to metals through many sources, including industrial wastewater discharge, mine drainage, roadways, and by the release of metals naturally occurring in near-surface rocks and sediments. Arizona Department of Environmental Quality (ADEQ) set standards to protect aquatic wildlife. Results for the following metals are compared to their appropriate standard: arsenic, cadmium, copper, lead, and zinc.

2008–2018 Summary

Most of the water samples tested for dissolved metals in the Santa Cruz River met the appropriate standards. The only samples with concentrations above the standard included a single sample for lead and several samples for cadmium. Cadmium is lethal to aquatic wildlife at relatively low

concentrations and comes from numerous human-produced sources such as electroplating processes, fertilizers, pesticides, coal combustion, and mine wastes. The cadmium standard was made more stringent in 2009, but in 2016 reverted back to a less stringent standard. Between 2008 and 2011, 11 (2009 standard) or 3 (2016 standard) samples tested for cadmium did not meet the standard. The treatment plant is not designed to remove metals, so ADEQ and the U.S. International Boundary and Water Commission worked with the cities of Nogales, Arizona and Nogales, Sonora to mitigate metals in the binational wastewater. These actions appear to have been successful as subsequent monitoring has not detected cadmium.

Although data are limited for Nogales Wash with samples collected from 2008–2012, only two samples had metals detected above the standard.

Nickel is not tracked in this assessment. However, a 2018 presentation by ADEQ suggests that nickel may be another contaminant to monitor and mitigate.

Values for dissolved metals in water samples tested throughout all water years
concentrations in micrograms/liter ($\mu\text{g/L}$), also known as parts per billion (ppb)

	Average (min–max)	Samples Above Standard	Average (min–max)	Samples Above Standard	Average Standard wildlife standards vary with water hardness
Arsenic	3.6 (3.1–4.9)	0	7.7 (3.8–17)	0	150 $\mu\text{g/L}$ ^a
Cadmium	3.9 ^b	1	1.4 (0.1–8.3)	11 or 3 ^c	0.8 or 3.7 $\mu\text{g/L}$ ^d
Copper	5.5 (1.9–18)	1	5.4 (2.9–11)	0	16 $\mu\text{g/L}$
Lead	ND	0	14 (0.04–40)	1	4.9 $\mu\text{g/L}$
Zinc	ND	0	42 (13–150)	0	211 $\mu\text{g/L}$
Nogales Wash ^e			Santa Cruz River		ND = Not detected

^aNot an average. Standard for arsenic does not vary with water hardness.

^bNot an average. Only one sample had a concentration detected in 2008 (which was above the standard). All others had concentrations of cadmium that were below detection level.

^cStandard changed in 2009 and 2016. Number of samples with concentrations above the standard varies; 11 with more stringent standard or 3 with less stringent standard. No samples were above the standard since 2011.

^dAverage standard for 2009 more stringent standard or average standard for 2016 less stringent standard.

^eSamples from Nogales Wash only available for 2008–2012 water years.



AQUATIC WILDLIFE: Fish


Fish can serve as effective longer-term indicators of river health because they live for several years and differ from species to species in their tolerance to pollution. Historically, the Santa Cruz River from Rio Rico to Amado supported several native fish species: Gila topminnow, desert sucker, Sonora sucker, and longfin dace. Long-term monitoring has shown a decline in the number of native fish species present. Three nonnative fish species—the Western mosquitofish, green sunfish, and largemouth bass—were found in Tumacácori National Historical Park between 2001 and 2002. There is no standard for diversity of fish. The results from the 2008 water year will serve as a baseline for monitoring change.

2008–2018 Summary

Annual fish surveys in autumn determine fish presence and species in Nogales Wash and at several locations

downstream of the treatment plant. In 2008, only longfin dace were found, and in very low numbers (2 individual fish caught). After the treatment plant upgrade, number of fish species has increased from one to four (two native and two non-native). The endangered Gila topminnow returned to the river in 2015 and the relative ease of capturing them during surveys suggests they are thriving. Conditions in the Amado reach have been occasionally dry at the time of survey and thus may no longer represent habitat as streamflow becomes more variable.

Though not impacted by the treatment plant upgrade, Nogales Wash has also seen an increase in fish species, most notably with the return of the Gila topminnow in 2018. Though not included in this summary, longfin dace were found in Nogales Wash in 2007. The reason why no fish were found during the baseline year is unknown.





AQUATIC WILDLIFE: Aquatic Invertebrates


Aquatic invertebrates break down organic materials and are important prey for fish and other species. They also differ in their tolerances to pollution. Chironomidae (midges) are pollution tolerant and found in high numbers even with low oxygen levels and high organic matter. Ephemeroptera (mayflies) have exposed gills on the outside of their body, making them very pollution sensitive. The percent of the invertebrate community comprised of Ephemeroptera taxa is commonly used to help track changes in water quality. Regardless of sensitivity to pollution, if a single species or group accounts for more than 50% of the community, this lack of diversity suggests a stream is impaired. Another way to look at diversity is simply looking at the total number of different invertebrate taxa found in the samples collected.

2008–2018 Summary

Information about the aquatic invertebrate community in the river is very limited. Our best data set is from Tumacácori National Historical Park in the Tubac Reach. Invertebrates

were sampled in 2008, and from 2012–2018 using a protocol that involved kick-net samples in riffles. Targeted riffle data of this kind are not available from 2009 and 2010. Other data collected with different methodology are included here to provide the most complete dataset. Data from 2011 are unavailable. The percent of the community dominated by a single group has decreased, but is still hovering around the 50 percent threshold. This suggests that the Tubac reach is under environmental stress. However, the percent of the aquatic community comprised of pollution-sensitive Ephemeroptera increased after the treatment plant upgrade, suggesting improvements in aquatic conditions. The number of taxa also has increased, providing further support of improved conditions. The park experienced dry conditions and more variable flow in 2013 and 2014, which may help explain some of the variability seen between 20013 and 2015. Further monitoring will be needed to determine if the aquatic invertebrate community will continue to improve.

Nogales Wash has no aquatic invertebrate data.



|| Treatment plant upgrades complete



GROUNDWATER: Maximum Depth

The interactions between riparian vegetation, surface water, and groundwater are important factors in overall riparian health. As with other southwestern rivers, cottonwoods and willows dominate the native riparian forests along the Santa Cruz River in this reach and depend on surface water and groundwater to meet their annual water requirements.


Several scientific studies have investigated the maximum depth to groundwater required to sustain mature cottonwood trees. Scientists estimate that the maximum depth to groundwater required to sustain mature Fremont cottonwoods ranges from 2.5 to 5.1 meters (approximately 8 to 16 feet).

We compare maximum depths to groundwater to the upper end of the cottonwood range, thus locations with maximum depths less than 5.1 meters meet the scientific standard. It is important to note that areas with greater maximum depths to groundwater support other types of riparian vegetation.

Depths to groundwater along Nogales Wash were not compiled for this summary.

2008–2018 Summary

Number of depth measurements over the years varied by well. Some wells were measured only once a year while others were measured daily. All depth to groundwater measurements are determined relative to areas that may support cottonwoods. The Tubac reach appears to have the most stable depth to groundwater measures. Rio Rico and Amado likely had increased recharge in recent years with shallower depths recorded. In Rio Rico, the wells with measurements below the standard are located near the beginning of the reach and represent wells closest to the release of effluent. In contrast, several wells in the Amado reach that were below the standard are located at the far end of the reach and represent wells that are furthest away from the release of effluent. The best conditions for supporting cottonwoods may be along the Tubac reach where geologic and hydrologic conditions are optimal and the recharge of effluent is possibly highest.






GROUNDWATER: January to June Decline

Variability in groundwater levels affects the growth and survival of riparian plants. There are several natural and human causes of groundwater variability. Naturally, groundwater levels decline after the winter rainy season ends and increase once the summer monsoons begin. Studies on the San Pedro River in Arizona indicate that cottonwood trees can withstand declines in groundwater levels up to 0.8 meters (2.6 feet) from January to June. Wells with groundwater levels that decline no more than 0.8 meters between January and June met this scientific standard.

2008–2018 Summary

The number of wells with enough data to calculate the change in depth to groundwater from January to June varied over the years. Some wells with less frequent monitoring did not have enough data or data were collected at the incorrect

time to determine the decline. As with maximum depth to groundwater, variability in depth to groundwater in the Tubac reach appears stable and the January to June standard was met at four well locations. Groundwater variability was less stable in the Rio Rico and Amado reaches, though data are limited. Rio Rico depths to groundwater varied between two wells. The well with less frequent monitoring is close to the Tubac reach and met the standard. The Rio Rico well with daily monitoring is near the effluent release and has greater fluctuations in water levels, often failing to meet the decline standard. Although both of the wells in the Amado reach met the standard, overall depth to groundwater may not support cottonwoods. Sonoran Institute launched a groundwater monitoring program in the Amado reach in 2018 that will help provide additional data on groundwater levels.





RIPARIAN VEGETATION: Forest and Woodland Cover

Riparian vegetation represents a small percentage of the land cover in the Santa Cruz watershed, but it provides important benefits to the region. Riparian vegetation's many ecosystem services include filtering contaminants from effluent-dominated water before it infiltrates into groundwater drinking supplies, slowing flood waters, reducing erosion potential along stream banks, increasing groundwater recharge, providing habitat for wildlife species, and providing recreational and spiritual enjoyment.

In 2006, Santa Cruz County, Friends of the Santa Cruz River, the University of Arizona, and the Sonoran Institute mapped the riparian vegetation along the Santa Cruz River from Rio Rico to Amado. Though there is no scientific standard quantifying the amount of riparian vegetation necessary to maintain a healthy river, a simplified version of the 2006 map establishes a baseline useful for measuring future changes in extent or composition of vegetation.

Types of vegetation and land use within the river's 100-year floodplain are divided into five categories: cottonwood forests and woodlands; mesquite forests and woodlands; other riparian vegetation; human infrastructure; and agriculture and pasture. Forests and woodlands have more than 10 percent cottonwood or mesquite tree cover. Other riparian vegetation includes areas not dominated by cottonwood or mesquite trees. Human infrastructure includes areas with

roads, railroads, and housing. Agriculture and pasture include crop fields and pasture land.

2008–2018 Summary

As of 2010, there were no major changes in vegetation type and landcover relative to the 2006 baseline. Only about 100 acres (1.5%) of the area within the 100-year floodplain had changes in vegetation type and landcover. In total, there were over 1,000 acres (16%) dominated by cottonwoods; nearly 1,400 acres (20%) dominated by mesquites; about 2,000 acres (31%) dominated by other riparian vegetation; over 500 acres (8%) used for human infrastructure; and 1,700 acres (25%) used for crops or pastures. Of the area in the 100-year floodplain, the Tubac reach had the largest percentage dominated by cottonwoods (26%), while the Rio Rico reach had the largest percentage dominated by mesquites (28%).

A detailed analysis of vegetation change between 2010 and 2018 has not been completed. However, casual field observations of locations in the Amado reach suggest that cottonwood forest and woodland cover in this area have suffered and trees have died or are showing significant die-back. With more variable flow extent, this is not surprising and we may see a shift to other riparian vegetation with increased distance from the treatment plant.

No vegetation analysis has been done for Nogales Wash.



LEARN MORE

Find all the *Living River* reports for Santa Cruz River at www.tiny.cc/livingriver.

ACKNOWLEDGEMENTS

Sonoran Institute Prepared this report with funding from Giles W. and Elise G. Mead Foundation, The William and Flora Hewlett Foundation, Tucson Electric Power, and community individuals.

The Sonoran Institute convened a Living River Technical Committee of ecology, hydrology, and wildlife experts to bring the best available science to bear on the development of the *Living River* health assessments.

The Technical Committee provided guidance by selecting and aggregating indicators of river health, identifying reference values or standards for evaluating and tracking changes in river conditions, and reviewing this report. The information presented in this report grew out of discussions involving these experts and represents the product of a collective effort; it does not reflect the opinions or viewpoints of any individual member of the technical team. The viewpoints and opinions expressed in the discussions of the group and captured in this report also do not reflect the opinions or viewpoints of the agencies, institutions, or organizations with whom the technical team members and external reviewers are associated or employed. Any errors or omissions contained herein are solely those of the Sonoran Institute.

Data sources: Arizona Department of Water Resources, Arizona Department of Environmental Quality, Friends of the Santa Cruz River, International Boundary and Water Commission, National Weather Service, National Park Service Sonoran Desert Network, Sonoran Institute, Tumacácori National Historical Park, and U.S. Geological Survey.

Production Credits:

Research, writing, and design - Claire A. Zugmeyer

Editing - Luke Cole

Map and illustrations on page 2, 3, and 18 respectively, icons, and design template - Terry Moody

Cover photos ©Bill Hatcher/Sonoran Institute, 2019



SONORAN
INSTITUTE

100 N. Stone Ave., Suite 400
Tucson, Arizona 85701

NON PROFIT
U.S. Postage
PAID
Permit #1454
Tucson, AZ 85701

SONORAN INSTITUTE has worked since our founding in 1990 to realize our vision that the Santa Cruz River, from Mexico to Marana, is a living, flowing river and the foundation of community health and prosperity.

The Sonoran Institute's mission is to connect people and communities with the natural resources that nourish and sustain them. We work at the nexus of commerce, community, and conservation to help people in the North American West build the communities they want to live in while preserving the values which brought them here. We envision a West where civil dialogue and collaboration are hallmarks of decision making, where people and wildlife live in harmony, and where clean water, air, and energy are assured.



Sonoran Institute



@sonoraninst



SonoranInstitute



WE WANT TO HEAR FROM YOU!

Help us map high-valued areas along the river by telling us where you visit the river: www.tiny.cc/scrmap

Learn more: SonoranInstitute.org



CleanWatts



Energy
CERTIFIED

Printed with 100% New Wind Energy